STANDARD F Tnl FIA performs quantitative measurement of Troponin I (Tnl) level in plasma and whole blood samples using fluorescence immunochromatography. Through the STANDARD F Analyzer, test results with high sensitivity and specificity can be obtained.


- Tnl is a cardiac marker for the diagnosis of myocardial infarction
- Automatic strip recognition by reading the information stored in 2D barcode
- Reagents can be stored at room temperature: 2-30°C / 36-86°F
- Plasma and Whole blood samples available



 $\epsilon$ 

Specification		
Intended use	Quantitative measurement of TnI level for diagnosis of AMI	
Specimen Volume	Whole Blood (EDTA), Serum	
Specimen type	100 µl	
Measurement Range	0.05 - 20 ng/mL	
Reference Range	< 0.05 ng/mL	
Testing Time	10 mins	
Storage Condition	2 - 30°C / 36 - 86°F	

## **TEST PROCEDURE**



Collect 100  $\mu$ l of sample with a fixed volume dropper (100  $\mu$ l).

Dispense collected sample into the extraction buffer tube. Then, discard the used fixed volume dropper (100 µl).

Mix sample and buffer 2-3 times with the disposable dropper (100  $\mu$ l). Then collect 100  $\mu$ l of sample mixture.

Apply the sample mixture into the sample well of the test device and immediately press the start button.

## **METHOD COMPARISON**

Reference method vs STANDARD F CK-MB FIA			
Correlation with ECLIA Method	y=0.9677x - 0.0069; R=0.9918; n=120		
CV%	QCL=8.8% / QCM=7.4% / QCH=7.1%		
Differ (%)	Within 15% (Serum) / Within 20% (Whole Blood)		

## **ORDERING INFORMATION**

Category	Product	Pack Size
Cardiovascular	STANDARD F Tnl FIA	20 Test
	STANDARD F Tnl Control	Lv1 x 10 / Lv2 x 10

